

**SUMMARY OF PRODUCT CHARACTERISTICS**  
**(SmPC)**

**FITZKING PROMETHAZINE INJECTION 50MG/2ML**

### 1.3 Product Information

#### 1.3.1 Summary of Product Characteristics (SmPC)

##### **1. Name of the medicinal product**

FITZKING PROMETHAZINE HYDROCHLORIDE INJECTION 50MG/2ML

##### **2. Qualitative and quantitative composition**

Each ampoule contains 25mg/ml of the active substance Promethazine hydrochloride.

Each 2ml ampoule contains 50mg of Promethazine hydrochloride.

For a full list of excipients, see section 6.1.

##### **3. Pharmaceutical form**

Solution for injection.

Promethazine hydrochloride injection is a clear colourless or almost colourless solution.

##### **4. Clinical particulars**

###### **4.1 Therapeutic indications**

As symptomatic treatment for allergic conditions of the upper respiratory tract and skin including allergic rhinitis, urticaria and anaphylactic reactions to drugs and foreign proteins.

Sedation and treatment of insomnia in adults.

As an adjunct in preoperative sedation in surgery and obstetrics.

As a paediatric sedative.

###### **4.2 Posology and method of administration**

Route of administration: Intramuscular or intravenous (after dilution)

The usual dose is 25 - 50 mg by deep intramuscular injection, or, in emergency, by slow intravenous injection after dilution of the 2.5% solution to 10 times its volume with water for injections immediately before use.

Maximum parenteral dose 100 mg.

Elderly: No specific dosage recommendations.

Children: 6.25 - 12.5 mg for children from 5 - 10 years by deep intramuscular injection. Not for use in children under 2 years of age (see section 4.3).

###### **4.3 Contraindications**

Promethazine hydrochloride should not be used in patients in coma or suffering from CNS depression of any cause.

Promethazine hydrochloride should not be given to patients with a known hypersensitivity to promethazine or to any of the excipients.

Promethazine is contraindicated for use in children less than two years of age because of the potential for fatal respiratory depression..

Promethazine hydrochloride should be avoided in patients taking monoamine oxidase inhibitors up to 14 days previously.

#### **4.4 Special warnings and precautions for use**

Promethazine hydrochloride may thicken or dry lung secretions and impair expectoration. It should therefore be used with caution in patients with asthma, bronchitis or bronchiectasis.

Use with care in patients with severe coronary artery disease, narrow angle glaucoma, epilepsy or hepatic and renal insufficiency.

Caution should be exercised in patients with bladder neck or pyloro-duodenal obstruction.

The use of promethazine should be avoided in children and adolescents with signs and symptoms suggestive of Reye's Syndrome.

Promethazine may mask the warning signs of ototoxicity caused by ototoxic drugs e.g. salicylates. It may also delay the early diagnosis of intestinal obstruction or raised intracranial pressure through the suppression of vomiting.

Intravenous injection should be performed with extreme care to avoid extravasation or inadvertent intra-arterial injection, which could lead to necrosis and peripheral gangrene. If a patient complains of pain during intravenous injection, stop the injection immediately, as this may be a sign of extravasation or inadvertent intra-arterial injection. Intramuscular injection must also be performed carefully to avoid inadvertent subcutaneous injection, which could lead to local necrosis.

Promethazine hydrochloride contains Sodium Sulphite and may rarely cause severe hypersensitivity reactions and bronchospasm.

#### **4.5 Interaction with other medicinal products and other forms of interaction**

Promethazine hydrochloride will enhance the action of any anticholinergic agent, tricyclic antidepressant, sedative or hypnotic. Alcohol should be avoided during treatment. Promethazine hydrochloride may cause hypotension, and dosage adjustment of antihypertensive therapy may therefore be required. Promethazine hydrochloride may lower the convulsive threshold, and dosage adjustment of anticonvulsant medication may therefore be required. Promethazine hydrochloride may interfere with immunological urine pregnancy tests to produce false-positive or false-negative results. Promethazine hydrochloride should be discontinued at least 72 hours before the start of skin tests as it may inhibit the cutaneous histamine response thus producing false-negative results. Promethazine hydrochloride injection may increase glucose tolerance.

#### **4.6 Fertility, pregnancy and lactation**

Promethazine hydrochloride injection should not be used in pregnancy unless the physician considers it essential. The use of Promethazine hydrochloride is not recommended in the 2 weeks prior to delivery in view of the risk of irritability and excitement in the neonate.

Available evidence suggests that the amount excreted in milk is insignificant. However, there are risks of neonatal irritability and excitement.

#### **4.7 Effects on ability to drive and use machines**

Ambulant patients receiving Promethazine hydrochloride for the first time should not be in control of vehicles or machinery for the first few days until it is established that they are not hypersensitive to the central nervous effects of the drug and do not suffer from disorientation, confusion or dizziness.

#### **4.8 Undesirable effects**

The following CIOMS frequency rating is used: Very common ( $\geq 1/10$ ); common ( $\geq 1/100$  to  $< 1/10$ ); uncommon ( $\geq 1/1000$  to  $< 1/100$ ); rare ( $\geq 1/10,000$  to  $\geq 1/1000$ ); very rare ( $< 1/10,000$ ), not known (cannot be estimated from the available data).

Side effects may be seen in a few patients: drowsiness, dizziness, restlessness, headaches, nightmares, tiredness, and disorientation. Anticholinergic side effects such as blurred vision, dry mouth and urinary retention occur occasionally. Newborn and premature infants are susceptible to the anticholinergic effects of promethazine, while other children may display paradoxical hyperexcitability. The elderly are particularly susceptible to the anticholinergic effects and confusion due to promethazine. Other side-effects include anorexia, gastric irritation, palpitations, hypotension, arrhythmias, extrapyramidal effects, muscle spasms and tic-like movements of the head and face. Jaundice and blood dyscrasias including haemolytic anaemia rarely occur. Very rare cases of allergic reactions, including urticaria, rash, pruritus and anaphylaxis, have been reported.

Photosensitive skin reactions have been reported; strong sunlight should be avoided during treatment.

#### **Reporting of suspected adverse reactions**

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard)

#### **4.9 Overdose**

Symptoms of severe overdosage are variable. They are characterised in children by various combinations of excitation, ataxia, incoordination, athetosis and hallucinations, while adults may become drowsy and lapse into coma. Convulsions may occur in both adults and children: coma or excitement may precede their occurrence. Cardiorespiratory depression is uncommon. If the patient is seen soon enough after ingestion, it should be possible to induce vomiting with ipecacuanha despite the antiemetic effect of promethazine; alternatively, gastric lavage may be used.

Treatment is otherwise supportive with attention to maintenance of adequate respiratory and circulatory status. Convulsions should be treated with diazepam or other suitable anticonvulsant.

## **5. Pharmacological properties**

### **5.1 Pharmacodynamic properties**

Pharmacotherapeutic group: Antihistamines for systemic use; Phenothiazine derivatives, ATC code: R06AD02

Potent, long acting, antihistamine with additional anti-emetic central sedative and anti-cholinergic properties

### **5.2 Pharmacokinetic properties**

Promethazine is slowly excreted via urine and bile. It is distributed widely in the body. It enters the brain and crosses the placenta. Phenothiazines pass into the milk at low concentrations.

### **5.3 Preclinical safety data**

No additional data of relevance to the prescriber.

## **6. Pharmaceutical particulars**

### **6.1 List of excipients**

Ascorbic acid

Anhydrous Sodium Sulphite

Sodium bisulphite

Water for injections

### **6.2 Incompatibilities**

Not applicable

### **6.3 Shelf life**

3 years

### **6.4 Special precautions for storage**

Store in the original carton in order to protect from light.

### **6.5 Nature and contents of container**

2 ml ampoules.

## **6.6 Special precautions for disposal and other handling**

Discoloured solutions should not be used.

## **7. Manufacturer**

JIANGSU HUAYANG PHARM. CO. LTD., NO. 12, CHANHGJIANG ROAD,  
SIYANG COUNTY, JIANGSU, CHINA

Fitzking Promethazine Hydrochloride injection 50mg/2ml