



1. Name of the medicinal product

CELICET 400 (Cefixime Tablets USP 400 mg)

2. Qualitative and quantitative composition

Cefixime Trihydrate USP Eq. to Cefixime Anhydrous ...400 mg

3. Pharmaceutical form

Film Coated Tablet.

4. Clinical particulars

4.1 Therapeutic indications

Uncomplicated Urinary Tract Infections

CELICET is indicated in the treatment of adults and pediatric patients six months of age or older with uncomplicated urinary tract infections caused by susceptible isolates of *Escherichia coli* and *Proteus mirabilis*.

Otitis Media

CELICET is indicated in the treatment of adults and pediatric patients six months of age or older with otitis media caused by susceptible isolates of *Haemophilus influenzae*, *Moraxella catarrhalis*, and *Streptococcus pyogenes*. (Efficacy for *Streptococcus pyogenes* in this organ system was studied in fewer than 10 infections.)

Pharyngitis And Tonsillitis

CELICET is indicated in the treatment of adults and pediatric patients six months of age or older with pharyngitis and tonsillitis caused by susceptible isolates of *Streptococcus pyogenes*.

Acute Exacerbations Of Chronic Bronchitis

CELICET is indicated in the treatment of adults and pediatric patients six months of age or older with acute exacerbations of chronic bronchitis caused by susceptible isolates of *Streptococcus pneumoniae* and *Haemophilus influenzae*.

Uncomplicated Gonorrhea (Cervical/Urethral)

CELICET is indicated in the treatment of adults and pediatric patients six months of age or older with uncomplicated gonorrhea (cervical/urethral) caused by susceptible isolates of *Neisseria gonorrhoeae* (penicillinase-and non-penicillinase-producing isolates).

4.2 Posology and method of administration

The usual course of treatment is 7 days. This may be continued for up to 14 days if required.

Posology

Adults and Children over 10 years or weighing more than 50 kg:



BAROQUE PHARMACEUTICALS PVT.LTD

The recommended dose is 200 – 400 mg daily according to the severity of infection, given either as a single dose or in two divided doses.

Children under 10 years:

CELICET Tablets 400 mg are not recommended for use in children under 10 years old. The safety and efficacy of cefixime has not been established in children less than 6 months.

Elderly:

Elderly patients may be given the same dose as recommended for adults. Renal function should be assessed, and dosage should be adjusted in severe renal impairment.

Renal impairment:

CELICET may be administered in the presence of impaired renal function. Normal dose and schedule may be given in patients with creatinine clearances of 20 ml/min or greater. In patients whose creatinine clearance is less than 20 ml/min, it is recommended that a dose of 400 mg once daily should not be exceeded. The dose and regimen for patients who are maintained on chronic ambulatory peritoneal dialysis or haemodialysis should follow the same recommendation as that for patients with creatinine clearances of less than 20 ml/min.

Method for administration

For oral administration. Absorption of CELICET is not significantly modified by the presence of food.

4.3 Contraindications

Hypersensitivity to cephalosporin antibiotics or to any of the excipients listed in section 6.1

4.4 Special warnings and precautions for use

Encephalopathy

Beta-lactams, including cefixime, predispose the patient to encephalopathy risk (which may include convulsions, confusion, impairment of consciousness, movement disorders), particularly in case of overdose or renal impairment.

Severe cutaneous adverse reactions

Severe cutaneous adverse reactions (SCARS) including toxic epidermal necrolysis (TEN), Stevens-Johnson syndrome (SJS) drug rash with eosinophilia and systemic symptoms (DRESS), and acute generalised exanthematous pustulosis (AGEP) have been reported in association with cefixime. Patients should be informed about the signs and symptoms of serious skin manifestations and monitored closely. Treatment should be discontinued at the first appearance of skin rash, mucosal lesions, or any other sign of skin hypersensitivity.

CELICET should be given with caution to patients who have shown hypersensitivity to other drugs.

Hypersensitivity to penicillins

As with other cephalosporins, cefixime should be given with caution to patients with a history of hypersensitivity to penicillin, as there is some evidence of partial cross-allergenicity between the penicillins and cephalosporins.



BAROQUE PHARMACEUTICALS PVT.LTD

Patients have had severe reactions (including anaphylaxis) to both classes of drugs. If an allergic effect occurs with CELICET, the drug should be discontinued and the patient treated with appropriate agents if necessary.

Haemolytic anaemia

Drug-induced haemolytic anaemia, including severe cases with a fatal outcome, has been described for cephalosporins (as a class). The recurrence of haemolytic anaemia after re-administration of cephalosporins in a patient with a history of cephalosporin (including cefixime) –associated haemolytic anaemia has also been reported.

Acute renal failure

As with other cephalosporins, cefixime may cause acute renal failure including tubulointerstitial nephritis as an underlying pathological condition. When acute renal failure occurs, cefixime should be discontinued and appropriate therapy and/or measures should be taken.

Renal impairment

CELICET should be administered with caution in patients with markedly impaired renal function (see section 4.2).

Paediatric use

Safety of cefixime in premature or newborn infant has not been established (see section 4.2).

Antibiotic-associated colitis

Treatment with broad spectrum antibiotics alters the normal flora of the colon and may permit overgrowth of clostridia. Studies indicate that a toxin produced by *Clostridium difficile* is a primary cause of antibiotic-associated diarrhoea. Pseudomembranous colitis is associated with the use of broad-spectrum antibiotics (including macrolides, semi-synthetic penicillins, lincosamides and cephalosporins); it is therefore important to consider its diagnosis in patients who develop diarrhoea in association with the use of antibiotics. Symptoms of pseudomembranous colitis may occur during or after antibiotic treatment.

Management of pseudomembranous colitis should include sigmoidoscopy, appropriate bacteriologic studies, fluids, electrolytes and protein supplementation. If the colitis does not improve after the drug has been discontinued, or if the symptoms are severe, oral vancomycin is the drug of choice for antibiotic-associated pseudomembranous colitis produced by *C. difficile*. Other causes of colitis should be excluded.

4.5 Interaction with other medicinal products and other forms of interaction

Anticoagulants

In common with other cephalosporins, increases in prothrombin times have been noted in a few patients. Care should therefore be taken in patients receiving anticoagulation therapy.

Cefixime should be administered with caution to patients receiving coumarin-type anticoagulants, e.g. warfarin potassium. Since cefixime may enhance effects of the anticoagulants, prolonged prothrombin time with or without bleeding may occur.



Other forms of interaction

A false positive reaction for glucose in the urine may occur with Benedict's or Fehling's solutions or with copper sulphate test tablets, but not with tests based on enzymatic glucose oxidase reactions.

A false positive direct Coombs test has been reported during treatment with cephalosporin antibiotics, therefore it should be recognised that a positive Coombs test may be due to the drug.

4.6 Fertility, pregnancy and lactation

Reproduction studies have been performed in mice and rats at doses up to 400 times the human dose and have revealed no evidence of impaired fertility or harm to the fetus due to cefixime. In the rabbit, at doses up to 4 times the human dose, there was no evidence of a teratogenic effect; there was a high incidence of abortion and maternal death which is an expected consequence of the known sensitivity of rabbits to antibiotic-induced changes in the population of the microflora of the intestine. There are no adequate and well-controlled studies in pregnant women. CELICET should therefore not be used in pregnancy or in nursing mothers unless considered essential by the physician.

4.7 Effects on ability to drive and use machines

In the case of side effects such as encephalopathy (which may include convulsion, confusion, impairment of consciousness, movement disorders), the patient should not operate machines or drive a vehicle.

4.8 Undesirable effects

CELICET is generally well tolerated. The majority of adverse reactions observed in clinical trials were mild and self-limiting in nature.

The following adverse reaction (Preferred term# or equivalent) will be considered listed:

Blood and lymphatic system disorders:	Eosinophilia Hypereosinophilia Agranulocytosis Leucopenia Neutropenia Granulocytopenia Haemolytic anaemia Thrombocytopenia Thrombocytosis
---------------------------------------	---



BAROQUE PHARMACEUTICALS PVT.LTD

Gastrointestinal disorders:	Abdominal pain Diarrhoea* Dyspepsia Nausea Vomiting Flatulence
Hepatobiliary disorders:	Jaundice
Infections and infestations:	Pseudomembranous colitis Vaginitis
Investigations:	Aspartate aminotransferase increased Alanine aminotransferase increased Blood bilirubin increased Blood urea increased Blood creatinine increased
Nervous system disorders:	Dizziness Headache Cases of convulsions have been reported with cephalosporins including cefixime (frequency not known)** Beta-lactams, including cefixime, predispose the patient to encephalopathy risk (which may include convulsions, confusion, impairment of consciousness, movement disorders), particularly in case of overdose or renal impairment (frequency not known)**
Respiratory, thoracic and mediastinal disorders:	Dyspnoea
Renal and urinary disorders:	Acute renal failure with tubulointerstitial nephritis (see section 4.4).
Immune system disorders:	Anaphylactic reaction Angio-oedema Serum sickness-like reaction
Skin and subcutaneous tissue disorders:	Drug rash with eosinophilia and systemic symptoms (DRESS) Erythema multiforme Stevens-Johnson syndrome Toxic epidermal necrolysis Urticaria Rash Pruritus Acute generalised exanthematous pustulosis(AGEP) (see section 4.4)
General disorders and administrative site conditions:	Drug Fever Arthralgia Pyrexia Face oedema Genital pruritus



BAROQUE PHARMACEUTICALS PVT.LTD

The above mentioned listed adverse reactions have been observed during clinical studies and/or during marketed use.

Preferred term in MedDRA (v.14.0)

*Diarrhoea has been more commonly associated with higher doses. Some cases of moderate to severe diarrhoea have been reported; this has occasionally warranted cessation of therapy. CELICET should be discontinued if marked diarrhoea occurs.

**Cannot be estimated from available data

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via Yellow Card Scheme at: www.mhra.gov.uk/yellowcard or search for MHRA Yellow Card in the Google Play or Apple App Store.

4.9 Overdose

Adverse events experienced in higher than recommended doses were similar to those seen at normal doses. In the event of overdosage general symptomatic and general supportive measures are indicated as required.

5. Pharmacological properties

5.1 Pharmacodynamic properties

5.1.1 Pharmacotherapeutic group: third generation cephalosporin

5.1.2 ATC code: J01DD08

Cefixime is an oral third generation cephalosporin which has marked in vitro bactericidal activity against a wide variety of Gram-positive and Gram-negative organisms.

Clinical efficacy has been demonstrated in infections caused by commonly occurring pathogens including *Streptococcus pneumoniae*, *Streptococcus pyogenes*, *Escherichia coli*, *Proteus mirabilis*, *Klebsiella* species, *Haemophilus influenzae* (beta-lactamase positive and negative), *Branhamella catarrhalis* (beta-lactamase positive and negative) and *Enterobacter* species. It is highly stable in the presence of beta-lactamase enzymes.

Most strains of enterococci (*Streptococcus faecalis*, group D Streptococci) and Staphylococci (including coagulase positive and negative strains and methicillin-resistant strains) are resistant to cefixime. In addition, most strains of *Pseudomonas*, *Bacteroides fragilis*, *Listeria monocytogenes* and *Clostridia* are resistant to cefixime.

5.2 Pharmacokinetic properties

The absolute oral bioavailability of cefixime is in the range of 22 – 54%. Absorption is not significantly modified by the presence of food. Cefixime may therefore be given without regard to meals.



BAROQUE PHARMACEUTICALS PVT.LTD

From *in vitro* studies, serum or urine concentrations of 1 mcg/mL or greater were considered to be adequate for most common pathogens against which cefixime is active. Typically, the peak serum levels following the recommended adult or paediatric doses are between 1.5 – 3 mcg/ml. Little or no accumulation of cefixime occurs following multiple dosing.

The pharmacokinetics of cefixime in healthy elderly (age > 64 years) and young volunteers (11 – 35) compared the administration of 400 mg doses once daily for 5 days. Mean C_{max} and AUC

values were slightly greater in the elderly. Elderly patients may be given the same dose as the general population.

Cefixime is predominantly eliminated as unchanged drug in the urine. Glomerular filtration is considered the predominant mechanism. Metabolites of cefixime have not been isolated from human serum or urine.

Serum protein binding is well characterised for human and animal sera; cefixime is almost exclusively bound to the albumin fraction, the mean free fraction being approximately 30%. Protein binding of cefixime is only concentration dependent in human serum at very high concentrations which are not seen following clinical dosing.

Transfer of ^{14}C -labelled cefixime from lactating rats to their nursing offspring through breast milk was quantitatively small (approximately 1.5% of the mothers' body content of cefixime in the pup). No data are available on secretion of cefixime in human breast milk. Placental transfer of cefixime was small in pregnant rats dosed with labelled cefixime.

5.3 Preclinical safety data

There are no pre-clinical data of relevance to the prescriber which are additional to that already included in other sections of the Summary of Product Characteristics.

6. Pharmaceutical particulars

6.1 List of excipients

Ingredients	Specification
Microcrystalline cellulose	USP/NF
Pregelatinised Starch	USP/NF
Calcium Hydrogen Phosphate Dihydrate	USP/NF
Magnesium Stearate	USP/NF
Hypromellose	USP/NF
Macrogol 6000	USP/NF
Titanium Dioxide	USP/NF
Purified Water	USP/NF



BAROQUE PHARMACEUTICALS PVT.LTD

6.2 Incompatibilities

There are no major incompatibilities.

6.3 Shelf life

36 months.

6.4 Special precautions for storage

Store at temperature not exceeding 30°C.

6.5 Nature and contents of container

10 Tablets of CELICET-400 are packed in ALU-ALU-blister, such 1 blister packed in monocarton along with leaflet and such 10 monocartons packed in an outer carton.

6.6 Special precautions for usage / preparation before use

Do not chewed or crushed, swallow whole with water as per directed by physician.

6.7 Special precautions for disposal and other handling

None.

7. Marketed by:

Greenfield Pharmaceutical Limited.
5, gunning Street, Fegge, Onitsha,
Anambra State, Nigeria.

8. Manufactured by:

Baroque Pharmaceuticals Pvt. Ltd.
Sokhada-388620, Ta-Khambhat
Dist-Anand (Gujarat), India.

9. Exported by:

AMN life science Pvt. Ltd.
150, Sahajanand Estate,
Sarkhej, Tal. City, Ahmedabad,
Gujarat , India.

10. Marketing authorization number(s):

NA

11. Date of first authorization/renewal of the authorization:



BAROQUE PHARMACEUTICALS PVT.LTD

NA

12. Date of revision of the text

January 2022