



MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)Ahlcon Parenterals (India) Limited

1. NAME OF THE MEDICINAL PRODUCT

Paracetamol Infusion 1.0% w/v

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each 100 mL contains

Paracetamol BP ----- 1000 mg.

Water for Injections BP-----q.s.

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Solution for infusion.

4. CLINICAL PARTICULARS**4.1 Therapeutic Indications**

Paracetamol Infusion 1.0% w/v is indicated for the short-term treatment of moderate pain, especially following surgery and for the short-term treatment of fever, when administration by intravenous route is clinically justified by an urgent need to treat pain or hyperthermia and/or when other routes of administration are not possible.

4.2 Posology and method of administration

Intravenous route.

The 100 ml is restricted to adults, adolescents and children weighing more than 33 kg.

Posology

Dosing based on patient weight (please see the dosing table here below)

Patient weight	Dose per administration	Volume per administration	Maximum volume of Paracetamol Infusion per administration based on upper weight limits of group (mL)**	Maximum Daily Dose ***
≤10 kg *	7.5 mg/kg	0.75 mL/kg	7.5mL	30 mg/kg
> 10 kg to ≤33kg	15 mg/kg	1.5mL/kg	49.5mL	60mg/kg not exceeding 2g
> 33 kg to ≤50kg	15 mg/kg	1.5mL/kg	75 mL	60mg/kg not exceeding 3g

MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

Patient weight	Dose per administration	Volume per administration	Maximum volume per Administration **	Maximum Daily Dose ***
>50kg with additional risk factors for hepatotoxicity	1g	100 mL	100 mL	3g
> 50 kg and no additional risk factors for hepatotoxicity	1g	100 mL	100 mL	4 g

* **Pre-term newborn infants:** No safety and efficacy data are available for pre-term newborn infants (see section 5.2).

** Patients weighing less will require smaller volumes.

The minimum interval between each administration must be at least 4 hours. No more than 4 doses to be given in 24 hours.

The minimum interval between each administration in patients with severe renal insufficiency must be at least 6 hours.

*** **Maximum daily dose:** The maximum daily dose as presented in the table above is for patients that are not receiving other paracetamol containing products and should be adjusted accordingly taking such products into account.

Severe renal insufficiency:

It is recommended, when giving paracetamol to patients with severe renal impairment (creatinine clearance ≤ 30 mL/min), to increase the minimum interval between each administration to 6 hours (See section 5.2). In adults with hepatocellular insufficiency, chronic alcoholism, chronic malnutrition (low reserves of hepatic glutathione), dehydration:

The maximum daily dose must not exceed 3 g (see section 4.4).



Method of administration

Take care when prescribing and administering Paracetamol Infusion 1.0% w/ to avoid dosing errors due to confusion between milligram (mg) and milliliter (mL), which could result in accidental overdose and death. Take care to ensure the proper dose is communicated and dispensed. When writing prescriptions, include both the total dose in mg and the total dose in volume.

The paracetamol solution is administered as a 15-minute intravenous infusion.

4.3 Contraindications

Paracetamol Infusion 1.0% w/v is contraindicated:

- in patients with hypersensitivity to paracetamol or to propacetamol hydrochloride (prodrug of paracetamol) or to one of the excipients.
- in cases of severe hepatocellular insufficiency.

MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

4.4 Special warnings and special precautions for use

Warnings

RISK OF MEDICATION ERRORS

Take care to avoid dosing errors due to confusion between milligram (mg) and milliliter (mL), which could result in accidental overdose and death (see section 4.2).

It is recommended to use a suitable analgesic oral treatment as soon as this administration route is possible.

In order to avoid the risk of overdose, check that other medicines administered do not contain either paracetamol or propacetamol.

Doses higher than the recommended entails risk for very serious liver damage. Clinical symptoms and signs of liver damage (including fulminant hepatitis, hepatic failure, cholestatic hepatitis, cytolytic hepatitis) are usually first seen after two days of drug

administration with a peak seen usually after 4 - 6 days. Treatment with antidote should be given as soon as possible (See section 4.9)



Precautions for use

Paracetamol should be used with caution in cases of:

- hepatocellular insufficiency,
- severe renal insufficiency (creatinine clearance ≤ 30 mL/min) (see sections 4.2 and 5.2),
- chronic alcoholism,
- chronic malnutrition (low reserves of hepatic glutathione),
- dehydration.

4.5 Interaction with other Medicinal Products and other forms of interaction

- Probenecid causes an almost 2-fold reduction in clearance of paracetamol by inhibiting its conjugation with glucuronic acid. A reduction of the paracetamol dose should be considered for concomitant treatment with probenecid,
- Salicylamide may prolong the elimination $t_{1/2}$ of paracetamol,
- Caution should be paid to the concomitant intake of enzyme-inducing substances (see section 4.9).

MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

- Concomitant use of paracetamol (4 g per day for at least 4 days) with oral anticoagulants may lead to slight variations of INR values. In this case, increased monitoring of INR values should be conducted during the period of concomitant use as well as for 1 week after paracetamol treatment has been discontinued

4.6 Pregnancy and lactation

Pregnancy

Clinical experience of intravenous administration of paracetamol is limited. However, epidemiological data from the use of oral therapeutic doses of paracetamol indicate no undesirable effects on the pregnancy or on the health of the foetus / newborn infant.



Prospective data on pregnancies exposed to overdoses did not show an increase in malformation risk.

Reproductive studies with the intravenous form of paracetamol have not been performed in animals. However, studies with the oral route did not show any malformation or foetotoxic effects.

Nevertheless, Paracetamol Infusion 1.0% w/v should only be used during pregnancy after a careful benefit-risk assessment. In this case, the recommended posology and duration must be strictly observed.

Lactation

After oral administration, paracetamol is excreted into breast milk in small quantities. No undesirable effects on nursing infants have been reported.

Consequently, Paracetamol Infusion 1.0% w/v may be used in breast-feeding women.

4.7 Effects on ability to drive and use machines Not relevant.

4.8 Undesirable effects

As all paracetamol products, adverse drug reactions are rare (>1/10000, <1/1000) or very rare (<1/10000), they are described below:

MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

Organ system	Rare >1/10000, <1/1000	Very Rare < 1/10,000
--------------	---------------------------	----------------------



General	Malaise	Hypersensitivity reaction
Cardiovascular	Hypotension	
Liver	Increased levels of hepatic transaminases	
Platelet/blood		
Platelet/blood		Thrombocytopenia, Leucopenia, Neutropenia.

Frequent adverse reactions at injection site have been reported during clinical trials (pain and burning sensation).

Very rare cases of hypersensitivity reactions ranging from simple skin rash or urticaria to anaphylactic shock have been reported and require discontinuation of treatment.

Cases of erythema, flushing, pruritus and tachycardia have been reported.

4.9 Overdose

There is a risk of liver injury (including fulminant hepatitis, hepatic failure, cholestatic hepatitis, cytolytic hepatitis), particularly in elderly subjects, in young children, in patients with liver disease, in cases of chronic alcoholism, in patients with chronic malnutrition and in patients receiving enzyme inducers. Overdosing may be fatal in these cases.

- Symptoms generally appear within the first 24 hours and comprise: nausea, vomiting, anorexia, pallor, abdominal pain. Overdose, 7.5 g or more of paracetamol in a single administration in adults and 140 mg/kg of body weight in a single administration in children, causes hepatic cytolysis likely to induce complete and irreversible necrosis, resulting in hepatocellular insufficiency, metabolic acidosis and encephalopathy which may lead to coma and death. Simultaneously, increased levels of hepatic transaminases (AST, ALT), lactate dehydrogenase and bilirubin are observed together with decreased prothrombin levels that may appear 12 to 48 hours after administration.

Clinical symptoms of liver damage are usually evident initially after two days, and reach a maximum after 4 to 6 days.

MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

Emergency measures

- Immediate hospitalisation.
- Before beginning treatment, take a tube of blood for plasma paracetamol assay, as soon as possible after the overdose.
- The treatment includes administration of the antidote, N-acetylcysteine (NAC), by the i.v. or oral route, if possible before the 10th hour. NAC can, however, give some degree of protection even after 10 hours, but in these cases prolonged treatment is given.
- Symptomatic treatment.
- Hepatic tests must be carried out at the beginning of treatment and repeated every 24 hours. In most cases hepatic transaminases return to normal in one to two weeks with full restitution of liver function. In very severe cases, however, liver transplantation may be necessary.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic Properties

Pharmacotherapeutic group: Other Analgesics and Antipyretics,

ATC code: N02BE01

The precise mechanism of the analgesic and antipyretic properties of paracetamol has yet to be established; it may involve central and peripheral actions.

Paracetamol Infusion 1.0% w/v provides onset of pain relief within 5 to 10 minutes after the start of administration. The peak analgesic effect is obtained in 1 hour and the duration of this effect is usually 4 to 6 hours.

Paracetamol Infusion 1.0% w/v reduces fever within 30 minutes after the start of administration with a duration of the antipyretic effect of at least 6 hours.



5.2 Pharmacokinetic properties

Adults:

Absorption:

Paracetamol pharmacokinetics is linear up to 2 g after single administration and after repeated administration during 24 hours.

MODULE 1; ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

The bioavailability of paracetamol following infusion of 500 mg and 1 g of Paracetamol Infusion 1.0% w/v is similar to that observed following infusion of 1 g and 2 g propacetamol (corresponding to 500 mg and 1 g paracetamol respectively). The maximal plasma concentration (C_{max}) of paracetamol observed at the end of 15 minutes intravenous infusion of 500 mg and 1 g of Paracetamol Infusion 1.0% w/v is about 15 µg/mL and 30 µg/mL respectively.

Distribution:

The volume of distribution of paracetamol is approximately 1 L/kg.

Paracetamol is not extensively bound to plasma proteins.

Following infusion of 1 g paracetamol, significant concentrations of paracetamol (about 1.5 µg/mL) were observed in the Cerebro Spinal Fluid as and from the 20th minute following infusion.

Metabolism:

Paracetamol is metabolised mainly in the liver following two major hepatic pathways: glucuronic acid conjugation and sulphuric acid conjugation. The latter route is rapidly saturable at doses that exceed the therapeutic doses. A small fraction (less than 4%) is metabolised by cytochrome P450 to a reactive intermediate (N-acetyl benzoquinone imine) which, under normal conditions of use, is rapidly detoxified by reduced glutathione

and eliminated in the urine after conjugation with cysteine and mercapturic acid. However, during massive overdosing, the quantity of this toxic metabolite is increased.



Elimination:

The metabolites of paracetamol are mainly excreted in the urine. 90% of the dose administered is excreted in 24 hours, mainly as glucuronide (60-80%) and sulphate (20-30%) conjugates. Less than 5% is eliminated unchanged. Plasma half-life is 2.7 hours and total body clearance is 18 L/h.

Neonates, infants and children

The pharmacokinetic parameters of paracetamol observed in infants and children are similar to those observed in adults, except for the plasma half-life that is slightly shorter

MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

(1.5 to 2 h) than in adults. In neonates, the plasma half-life is longer than in infants i.e. around 3.5 hours. Neonates, infants and children up to 10 years excrete significantly less glucuronide and more sulphate conjugates than adults.

Table. Age related pharmacokinetic values (standardized clearance, *CL_{std}/F_{oral} (l.h⁻¹ 70 kg⁻¹), are presented below.

Age	Weight (kg)	CL _{std} /F _{oral} (l.h ⁻¹ 70 kg ⁻¹),
40 weeks PCA	3.3	6
3 months PNA	7.5	5.9
6 months PNA	10	8.8
1 year PNA	12	11.1 13.6
2 years PNA	20	15.6 16.3
5 years PNA	25	16.3
8 years PNA		

*CL_{std} is the population estimate for CL

Special populations:

Renal insufficiency



In cases of severe renal impairment (creatinine clearance 10-30 mL/min), the elimination of paracetamol is slightly delayed, the elimination half-life ranging from 2 to 5.3 hours. For the glucuronide and sulphate conjugates, the elimination rate is 3 times slower in subjects with severe renal impairment than in healthy subjects. Therefore, it is recommended, when giving paracetamol to patients with severe renal impairment

(creatinine clearance \leq 30 mL/min), to increase the minimum interval between each administration to 6 hours (see section 4.2. Posology and method of administration).

Elderly subjects

The pharmacokinetics and the metabolism of paracetamol are not modified in elderly subjects. No dose adjustment is required in this population.

5.3 Preclinical safety data

Preclinical data reveal no special hazard for humans beyond the information included in other sections of the SmPC.

Studies on local tolerance of Paracetamol Infusion in rats and rabbits showed good tolerability. Absence of delayed contact hypersensitivity has been tested in guinea pigs.

MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Mannitol

Disodium Hydrogen Phosphate (Anhydrous Disodium Hydrogen Phosphate)

Sodium Hydroxide

Hydrochloric Acid

Water for injections

6.2 Incompatibilities

Paracetamol Infusion 1.0% w/v should not be mixed with other medicinal products.



6.3 Shelf life

24 Months.

6.4 Special precautions for storage

Do not store above 30°C. Protect from light. Do not freeze.

Keep out of the sight and reach of Children.

6.5 Nature and contents of container 100 mL LDPE Bottle.

6.6 Special precautions for disposal and other handling

Before administration, the product should be visually inspected for any particulate matter and discoloration. For single use only. Any unused solution should be discarded.

7. MARKETING AUTHORISATION HOLDER

Mark Pharmaceuticals Limited

8. MARKETING AUTHORISATION NUMBER(s)

B4-4012

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

.....

10. DATE OF REVISION OF THE TEXT

.....

MODULE 1: ADMINISTRATIVE AND PRODUCT INFORMATION

Paracetamol Infusion 1000 mg/100 mL (1.0% w/v)

Ahlcon Parenterals (India) Limited

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

.....

10. DATE OF REVISION OF THE TEXT

.....
