

14. ERYFAST 500 TABLETS (ERYTHROMYCIN 500MG)
15. P-ESPEN TABLETS (IBUPROFEN 200MG, PARACETAMOL 325MG)
16. XANAP 50 TABLETS (DICLOFENAC POTASSIUM USP 50MG)
17. XANAP 100 TABLETS (DICLOFENAC POTASSIUM USP 100MG)
18. XANAP FORTE TABLETS (PARACETAMOL & DICLOFENAC POTASSIUM TABLETS)
19. (COLDRILIF CAPSULES (CHLORPHENIRAMINE MALEATE & PHENYLEPHRINE HYDROCHLORIDE CAPSULES)
20. ASHMATAC-F TABLETS (SALBUTAMOL & THEOPHYLLINE TABLETS)
21. LILYAPETI TABLETS (CYPROHEPTADINE TABLETS BP) 0
22. CHYMOGIN TABLETS (TRYPSIN/CHYMOTRYPSIN)
23. ZONAGIN TABLETS (ACETAMINOPHEN & CAFFEINE TABLETS USP)
24. FAMTER TABLETS (ARTEMETHER 20MG, LUMEFANTRINE 120MG) A4-0801
25. PARANOVO CAPSULES PARACETAMOL & CAFFEINE CAPSULES
26. SPEVIN CAPSULES (PIROXICAM USP 20MG)
27. 27. OMECET PLUS (CLARITHROMYCIN, OMEPRAZOLE, TINIDAZOLE COMBIKIT)
28. 28. LINCODON LINCOMYCIN HYDROCHLORIDE CAPSULES BP)
29. 29 DIFLAZON (50MG FLUCONAZOLE 50MG)

Please note that the inspection of the site will be carried out at a later date after the COVID -19 pandemic and that any failure to meet minimum NAFDAC GMP requirements at the time will invalidate this approval. Also, in view of the above you are required to give your commitment to provide a seamless involvement in processing of staff visas when the time comes, otherwise the approval shall be reversed.

You may reach out to Director Drug Registration and Regulatory Affairs Directorate for further processing of your application.

Thank you,



Mrs. Ijeoma U. Nwankwo
Director (DER)
For: Director General (NAFDAC)



National Agency for Food & Drug Administration & Control (NAFDAC)

Registration & Regulatory Affairs (R & R) Directorate

SUMMARY OF PRODUCT CHARACTERISTICS (SmPC)

1. Name of the medicinal product**Fanmet Tablet-** (Dihydroartemisinin 40 + Piperaquine phosphate 320 Tablets)**2. Qualitative and quantitative composition**

Sr. No.	Ingredients	Spec .	Unit Formula (mg)	Batch Formula (kg)
1	Dihydroartemisinin	IH	40.000	17.600* (#)
2	Piperaquine Phosphate	IH	320.000	140.800*
3	Mannitol	IH	18.000	7.92
4	Lactose	IH	45.500	20.020**
5	Sodium Lauryl Sulphate	BP	5.000	2.200
6	Carmellose Sodium	BP	17.500	7.700
7	PVP K-30	BP	11.500	5.060
8	Isopropyl Alcohol	BP	q.s.	75.000
9	Purified Talc	BP	7.000	3.080
10	Carmellose Sodium	BP	20.500	9.020
11	Colloidal Anhydrous Silica (Aerosil)	BP	8.000	3.520
12	Magnesium Stearate	BP	7.000	3.080
Weigh of Compressed Tablet			500.000	
Coating				
13	Insta Coat Universal (IC-U-1308 - White)	IH	9.351	4.937***
14	Indigo Carmine Supra	IH	0.063	0.028
15	Methylene dichloride	BP	q.s.	61.060
16	Isopropyl Alcohol	BP	q.s.	32.840
Weigh of Coated Tablet			509.414	

3. Pharmaceutical form

Film-coated tablet (tablet).

Description: Blue Circular shape, biconvex, film coated tablet having embossing of “FANMET” on one side and “MAXHEAL” on other side

White oblong biconvex film-coated tablet (dimension 16x8mm / thickness 5.5mm) with a break-line and marked on one side with two “σ” letters.

The tablet can be divided into equal doses.

4. Clinical particulars

4.1 Therapeutic indications

Eurartesim is indicated for the treatment of uncomplicated *Plasmodium falciparum* malaria in adults, adolescents, children and infants 6 months and over and weighing 5 kg or more.

Consideration should be given to official guidance on the appropriate use of antimalarial medicinal products.

4.2 Posology and method of administration

Posology

Eurartesim should be administered over three consecutive days for a total of three doses taken at the same time each day.

Dosing should be based on body weight as shown in the table below.

Body weight (kg)	Daily dose (mg)		Tablet strength and number of tablets per dose
	PQP	Artemimol	
5 to <7	80	10	½ x 160 mg / 20 mg tablet
7 to <13	160	20	1 x 160 mg / 20 mg tablet
13 to <24	320	40	1 x 320 mg / 40 mg tablet
24 to <36	640	80	2 x 320 mg / 40 mg tablets
36 to <75	960	120	3 x 320 mg / 40 mg tablets
75 to 100	1,280	160	4 x 320 mg / 40 mg tablets
>100	There are no data on which to base a dose recommendation in patients weighing >100 kg.		

If a patient vomits within 30 minutes of taking Eurartesim, the whole dose should be re-administered; if a patient vomits within 30-60 minutes, half the dose should be re-administered. Re-dosing with Eurartesim should not be attempted more than once. If the second dose is vomited, alternative antimalarial therapy should be instituted.

If a dose is missed, it should be taken as soon as realised and then the recommended regimen continued until the full course of treatment has been completed.

There is no data on a second course of treatment.

No more than two courses of Eurartesim may be given within a 12 month period (see sections 4.4 and 5.3).

A second course of Eurartesim should not be given within 2 months after the first course due to the long elimination half-life of piperazine (see sections 4.4 and 5.2).

Special populations

Elderly

Clinical studies of Eurartesim did not include patients aged 65 years and over, therefore no dosing recommendation can be made. Considering the possibility of age-associated decrease in hepatic and renal function, as well as a potential for heart disorders (see sections 4.3 and 4.4), caution should be exercised when administering the product to the elderly.

Hepatic and renal impairment

Eurartesim has not been evaluated in subjects with moderate or severe renal or hepatic insufficiency. Therefore, caution is advised when administering Eurartesim to these patients (see section 4.4).

Paediatric population

The safety and efficacy of Eurartesim in infants aged less than 6 months and in children weighing less than 5 kg has not been established. No data are available for these paediatric subsets.

Method of administration

Eurartesim should be taken orally with water and without food.
Each dose should be taken no less than 3 hours after the last food intake.
No food should be taken within 3 hours after each dose.

For patients unable to swallow the tablets, such as infants and young children, Eurartesim may be crushed and mixed with water. The mixture should be used immediately after preparation.

4.3 Contraindications

- Hypersensitivity to the active substances or to any of the excipients listed in section 6.1.
- Severe malaria according to WHO definition.
- Family history of sudden death or of congenital prolongation of the QTc interval.
- Known congenital prolongation of the QTc-interval or any clinical condition known to prolong the QTc interval.
- History of symptomatic cardiac arrhythmias or with clinically relevant bradycardia.
- Any predisposing cardiac conditions for arrhythmia such as severe hypertension, left ventricular hypertrophy (including hypertrophic cardiomyopathy) or congestive cardiac failure accompanied by reduced left ventricle ejection fraction.
- Electrolyte disturbances, particularly hypokalaemia, hypocalcaemia or hypomagnesaemia.
- Taking medicinal products that are known to prolong the QTc interval. These include (but are not limited to):
 - Antiarrhythmics (e.g. amiodarone, disopyramide, dofetilide, ibutilide, procainamide, quinidine, hydroquinidine, sotalol).
 - Neuroleptics (e.g. phenothiazines, sertindole, sultopride, chlorpromazine, haloperidol, mesoridazine, pimozide, or thioridazine), antidepressive medicinal products.
 - Certain antimicrobial medicinal products, including medicinal products of the following classes:
 - macrolides (e.g. erythromycin, clarithromycin),
 - fluoroquinolones (e.g. moxifloxacin, sparfloxacin),
 - imidazole and triazole antifungal medicinal products,
 - and also pentamidine and saquinavir.
 - Certain non-sedating antihistamines (e.g. terfenadine, astemizole, mizolastine).
 - Cisapride, droperidol, domperidone, bepridil, diphemanil, probucol, levomethadyl, methadone, vinca alkaloids, arsenic trioxide.

- Recent treatment with medicinal products known to prolong the QTc interval that may still be circulating at the time that Eurartesim is commenced (e.g. mefloquine, halofantrine, lumefantrine, chloroquine, quinine and other antimalarial medicinal products) taking into account their elimination half-life.

4.4 Special warnings and precautions for use

Eurartesim should not be used to treat severe falciparum malaria (see section 4.3) and, due to insufficient data, should not be used to treat malaria due to *Plasmodium vivax*, *Plasmodium malariae* or *Plasmodium ovale*.

The long half-life of piperazine (about 22 days) should be kept in mind in the event that another anti-malarial agent is started due to treatment failure or a new malaria infection (see below and sections 4.3 and 4.5).

Piperazine is a mild inhibitor of CYP3A4. Caution is recommended when co-administering Eurartesim with medicinal products exhibiting variable patterns of inhibition, induction or competition for CYP3A4 as the therapeutic and/or toxic effects of some co-administered medicinal products could be altered.

Piperazine is also a substrate of CYP3A4. A moderate increase of piperazine plasma concentrations (<2-fold) was observed when co-administered with strong CYP3A4 inhibitors, resulting in a potential exacerbation of the effect on QTc prolongation (see section 4.5).

Exposure to piperazine may also be increased when co-administered with mild or moderate CYP3A4-inhibitors (e.g. oral contraceptives). Therefore, caution should be applied when co-administering Eurartesim with any CYP3A4-inhibitor and ECG monitoring should be considered.

Due to the lack of multiple dose PK data for piperazine, administration of any strong CYP3A4-inhibitors should be discouraged after initiation (i.e. the first dose) of Eurartesim (see sections 4.5 and 5.2).

Eurartesim should not be used during pregnancy in situations where other suitable and effective antimalarials are available (see section 4.6).

In the absence of carcinogenicity study data, and due to lack of clinical experience with repeated courses of treatment in humans, no more than two courses of Eurartesim should be given in a 12-month period (see sections 4.2 and 5.3).

Effects on cardiac repolarization

In clinical trials with Eurartesim limited ECGs were obtained during treatment. These showed that QTc prolongation occurred more frequently and to a larger extent in association with Eurartesim therapy than with the comparators (see section 5.1 for details of the comparators). Analysis of cardiac adverse events in clinical trials showed that these were reported more frequently in Eurartesim treated patients than in those treated with comparator antimalarial (see section 4.8). Before the third dose of Eurartesim, in one of the two Phase III studies 3/767 patients (0.4%) were reported to have a QTcF value of >500 ms versus none in the comparator group.

The potential for Eurartesim to prolong the QTc interval was investigated in parallel groups of healthy volunteers who took each dose with high (~1000 Kcal) or low (~400 Kcal) fat/calorie meals or in fasting conditions. Compared to placebo, the maximum mean increases in QTcF on Day 3 of dosing with Eurartesim were 45.2, 35.5 and 21.0 msec under respective dosing conditions. The QTcF prolongation observed under fasting conditions lasted between 4 and 11 hours after the last dose was administered on Day 3. The mean QTcF prolongation compared to placebo decreased to 11.8 msec at 24 hours and to 7.5 msec at 48 hours. No healthy subject dosed in fasting conditions showed a QTcF greater than 480 msec or an increase over baseline greater than 60 msec. The number of subjects with QTcF greater than 480 msec after dosing with low fat meals was 3/64, while 10/64 had QTcF values over this threshold after dosing with high fat meals. No subject had a QTcF value greater than 500 msec in any of the dosing conditions.

An ECG should be obtained as early as possible during treatment with Eurartesim and ECG monitoring should be applied in patients who may have a higher risk of developing arrhythmia in association with QTc prolongation (see below).

When clinically appropriate, consideration should be given to obtaining an ECG from all patients before the last of the three daily doses is taken and approximately 4-6 hours after the last dose, since the risk of QTc interval prolongation may be greatest during this period (see section 5.2). QTc intervals of more than 500 ms are associated with a pronounced risk for potentially life-threatening ventricular tachyarrhythmias. Therefore, ECG monitoring during the following 24-48 hours should be applied for patients found to have a prolongation to this extent. These patients should not receive another dose of Eurartesim and alternative antimalarial therapy should be instituted.

Compared to adult males, female patients and elderly patients have longer QTc intervals. Therefore, they may be more sensitive to the effects of QTc-prolonging medications such as Eurartesim so that special caution is required.

Paediatric population

Special precaution is advised in young children when vomiting, as they are likely to develop electrolyte disturbances. These may increase the QTc-prolonging effect of Eurartesim (see section 4.3).

Hepatic and renal impairment

Eurartesim has not been evaluated in patients with moderate or severe renal or hepatic insufficiency (see section 4.2). Due to the potential for higher plasma concentrations of piperazine to occur, caution is advised if Eurartesim is administered to patients with jaundice and/or with moderate or severe renal or hepatic insufficiency, and ECG and blood potassium monitoring are advised.

4.5 Interaction with other medicinal products and other forms of interaction

Eurartesim is contraindicated in patients already taking other medicinal products that are known to prolong the QTc interval due to the risk of a pharmacodynamic interaction leading to an additive effect on the QTc interval (see sections 4.3 and 4.4).

A limited number of drug-drug pharmacokinetic interaction studies with Eurartesim have been performed in healthy adult subjects. Therefore the assessment of the potential for drug-drug interactions to occur is based on either *in vivo* or *in vitro* studies.

Effect of Eurartesim on co-administered medicinal products

Piperazine is metabolised by, and is an inhibitor of CYP3A4. The concurrent administration of oral Eurartesim with 7.5 mg oral midazolam, a CYP3A4 probe substrate, led to a modest increase (≤ 2 -fold) in midazolam and its metabolites exposures in healthy adult subjects. This inhibitory effect was no longer evident one week after last administration of Eurartesim. Therefore, particular attention should be paid when medicinal products that have a narrow therapeutic index (e.g. antiretroviral medicinal products and cyclosporine) are co-administered with Eurartesim.

From *in vitro* data, piperazine undergoes a low level of metabolism by CYP2C19, and is also an inhibitor of this enzyme. There is the potential for reducing the rate of metabolism of other substrates of this enzyme, such as omeprazole, with consequent increase of their plasma concentration, and therefore, of their toxicity.

Piperazine has the potential to increase the rate of metabolism for CYP2E1 substrates resulting in a decrease in the plasma concentrations of substrates such as paracetamol or theophylline, and the anaesthetic gases enflurane, halothane and isoflurane. The main consequence of this interaction could be a reduction of efficacy of the co-administered medicinal products.

Artemimol administration may result in a slight decrease in CYP1A2 activity. Caution is therefore, advised when Eurartesim is administered concomitantly with medicinal products metabolised by

this enzyme that have a narrow therapeutic index, such as theophylline. Any effects are unlikely to persist beyond 24 hours after the last intake of arteminol.

Effect of co-administered medicinal products on Eurartesim

Piperaquine is metabolised by CYP3A4 *in vitro*. The concurrent administration of a single dose of oral clarithromycin, (a strong CYP3A4 inhibitor probe) with a single dose of oral Eurartesim led to a modest increase (≤ 2 -fold) in piperaquine exposure in healthy adult subjects. This increase in exposure to the antimalarial combination may result in an exacerbation of the effect on QTc (see section 4.4). Therefore, particular caution is required if Eurartesim is administered to patients taking potent CYP3A4 inhibitors (e.g. some protease inhibitors [amprenavir, atazanavir, indinavir, nelfinavir, ritonavir], nefazodone or verapamil), and ECG monitoring should be considered due to the risk of higher plasma concentrations of piperaquine (see section 4.4).

Enzyme inducing medicinal products such as rifampicin, carbamazepine, phenytoin, phenobarbital, St. John's wort (*Hypericum perforatum*) are likely to lead to reduced piperaquine plasma concentrations. The concentration of arteminol may also be reduced. Concomitant treatment with such medicinal products is not recommended.

Paediatric population

Drug-drug interaction studies have only been performed in adults. The extent of interactions in the paediatric population is not known. The above mentioned interactions for adults and the warnings in section 4.4 should be taken into account for the paediatric population.

Oral contraceptives

When co-administered to healthy women, Eurartesim exerted only a minimum effect on an estrogen/progestinic combination oral contraceptive treatment increasing the ethynilestradiol rate of absorption (expressed by geometric mean C_{max}) of about 28% but not significantly changing the exposure to ethynilestradiol and levonorgestrel and not influencing contraception activity as demonstrated by the similar plasma concentrations of follicle stimulating hormone (FSH), luteinizing hormone (LH) and progesterone observed after oral contraceptive treatment with or without concomitant Eurartesim administration.

Food interaction

Absorption of piperaquine is increased in the presence of fatty food (see sections 4.4 and 5.2) which may increase its effect on QTc interval. Therefore, Eurartesim should be taken with water only as described in section 4.2. Eurartesim should not be taken with grapefruit juice as it is likely to lead to increased piperaquine plasma concentrations.

4.6 Pregnancy and lactation

Pregnancy

There are insufficient data on the use of arteminol and piperaquine in pregnant women. Based on animal data, Eurartesim is suspected to cause serious birth defects when administered during the first trimester of pregnancy (see sections 4.4 and 5.3). Reproductive studies with artemisinin derivatives have demonstrated teratogenic potential with an increased risk during early gestation (see section 5.3). Piperaquine was not teratogenic in the rat or rabbit. In perinatal and postnatal studies in rats, piperaquine was associated with delivery complications. However, there was no delay in neonatal development following exposure *in utero* or via milk.

Eurartesim should not be used during pregnancy in situations where other suitable and effective anti-malarials are available (see section 4.4).

Breast-feeding

Animal data suggest excretion of piperaquine into breast milk but no data are available in humans. Women taking Eurartesim should not breast-feed during their treatment.

Fertility

There are no specific data relating to the effects of piperazine on fertility, however, to date no adverse events have been reported during clinical use. Moreover, data obtained in animal studies show that fertility is unaffected by arteminol in both females and males.

4.7 Effects on ability to drive and use machines

Adverse event data collected in clinical trials suggest that Eurartesim has no influence on the ability to drive and operate machines once the patient has recovered from the acute infection.

4.8 Undesirable effects

Summary of the safety profile

The safety of Eurartesim has been evaluated in two phase III open-label studies involving 1,239 paediatric patients up to 18 years and 566 adult patients >18 years treated with Eurartesim.

In a randomized trial in which 767 adults and children with uncomplicated *P. falciparum* malaria were exposed to Eurartesim, 25% of subjects were judged to have experienced an adverse drug reaction (ADR). No single type of ADR occurred at an incidence of $\geq 5\%$. The most frequent ADRs observed at an incidence $\geq 1.0\%$ were: Headache (3.9%), Electrocardiogram QTc Prolonged (3.4%), *P. falciparum* infection (3.0%), Anaemia (2.8%), Eosinophilia (1.7%), Haemoglobin decreased (1.7%), Sinus tachycardia (1.7%), Asthenia (1.6%), Haematocrit [decreased] (1.6%), Pyrexia (1.5%), Red Blood Cell Count decreased (1.4%). A total of 6 (0.8%) subjects had serious ADRs in the study.

In a second randomized trial, 1,038 children, aged between 6 months and 5 years, were exposed to Eurartesim and 71% were judged to have experienced an ADR. The following ADRs were observed at an incidence of $\geq 5.0\%$: Cough (32%), Pyrexia (22.4%), Influenza (16.0%), *P. falciparum* infection (14.1%), Diarrhoea (9.4%), Vomiting (5.5%) and Anorexia (5.2%). A total of 15 (1.5%) subjects had serious ADRs in the study.

Tabulated list of adverse reactions

In the tables below, ADRs are listed under system organ class (SOC), and ranked by headings of frequency. Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness, using the following convention: Very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$), not known (cannot be estimated from the available data). The table in this section is for adult patients only. A corresponding table for paediatric patients is presented in the specific section below.

Frequency of ADRs in adult patients participating in clinical studies with Eurartesim:

SOC	Very Common	Common	Uncommon
Infections and infestations		<i>P. falciparum</i> infection	Respiratory tract infection Influenza
Blood and lymphatic system disorders		Anaemia	
Metabolism and nutrition disorders			Anorexia
Nervous system disorders		Headache	Convulsion Dizziness
Cardiac disorders		QTc prolonged Tachycardia	Cardiac conduction disorders Sinus arrhythmias Bradycardia

Respiratory, thoracic and mediastinal disorders			Cough
Gastrointestinal disorders			Vomiting Diarrhoea Nausea Abdominal pain
Hepatobiliary disorders			Hepatitis Hepatomegaly Abnormal liver function tests
Skin and subcutaneous Tissue disorders			Pruritis
Musculoskeletal and connective tissue disorders			Arthralgia Myalgia
General disorders and administration site conditions		Asthenia Pyrexia	

Description of selected adverse reactions

The ADRs noted for Eurartesim were generally mild in severity, and the majority were non-serious. Reactions such as cough, pyrexia, headache, *P. falciparum* infection, anaemia, asthenia, anorexia and the observed changes in blood cell parameters are consistent with those expected in patients with acute malaria. The effect on prolongation of the QTc interval was observed on Day 2, and had resolved by Day 7 (the next time point at which ECGs were performed).

Paediatric population

A tabular overview of the frequency of the ADRs in paediatric patients is given below. The majority of paediatric experience is derived from African children aged 6 months to 5 years.

Frequency of ADRs in paediatric patients participating in clinical studies with Eurartesim:

SOC	Very Common	Common	Uncommon
Infections and infestations	Influenza <i>P. falciparum</i> infection	Respiratory tract infection Ear infection	
Blood and lymphatic system disorders		Thrombocytopenia Leukopenias/neutropenia Leucocytoses NEC Anaemia	Thrombocythaemia Splenomegaly Lymphadenopathy Hypochromasia
Metabolism and nutrition disorders		Anorexia	
Nervous system disorders			Convulsion Headache
Eye disorders		Conjunctivitis	
Cardiac disorders		QT/QTc prolonged Heart rate irregular	Cardiac conduction disorders Cardiac murmur

Respiratory, thoracic and mediastinal disorders	Cough		Rhinorrhoea Epistaxis
Gastrointestinal disorders		Vomiting Diarrhoea Abdominal pain	Stomatitis Nausea
Hepatobiliary disorders			Hepatitis Hepatomegaly Abnormal liver function tests Jaundice
Skin and subcutaneous tissue disorders		Dermatitis Rash	Acanthosis Pruritis
Musculoskeletal and connective tissue disorders			Arthralgia
General disorders and administration site conditions	Pyrexia	Asthenia	

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the the Yellow Card Scheme at the website: www.mhra.gov.uk/yellowcard.

4.9 Overdose

In clinical trials, nine patients received double the cumulative intended dose of Eurartesim. The safety profile of these patients did not differ from that of patients receiving the recommended dose, with no patient reporting SAEs.

In cases of suspected overdose, symptomatic and supportive therapy should be given as appropriate, including ECG monitoring because of the possibility of QTc interval prolongation (see section 4.4)

5. Pharmacological properties

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antiprotozoals, antimalarials, artemisinin and derivatives, combinations, ATC code: P01BF05.

Pharmacodynamic effects

Artemimol is able to reach high concentrations within the parasitized erythrocytes. Its endoperoxide bridge is thought to be essential for its antimalarial activity, causing free-radical damage to parasite membrane systems including:

- Inhibition of *falciparum* sarcoplasmic-endoplasmic reticulum calcium ATPase,
- Interference with mitochondrial electron transport
- Interference with parasite transport proteins
- Disruption of parasite mitochondrial function

The exact mechanism of action of piperazine is unknown, but it likely mirrors that of chloroquine, a close structural analogue. Chloroquine binds to toxic haeme (derived from the patient's

haemoglobin) within the malaria parasite, preventing its detoxification via a polymerisation step. Piperaquine is a bisquinoline, and this class has shown good antimalarial activity against chloroquine-resistant *Plasmodium* strains *in vitro*. The bulky bisquinolone structure may be important for activity against chloroquine-resistant strains, and may act through the following mechanisms:

- Inhibition of the transporters that efflux chloroquine from the parasite food vacuole
- Inhibition of haem-digestion pathway in the parasite food vacuole.

Resistance to piperaquine (when used as monotherapy) has been reported.

The efficacy and safety of Eurartesim have been assessed in two large randomised, open-label clinical trials:

Study DM040010 was conducted in Asian adult and paediatric patients with uncomplicated *P. falciparum* malaria. Eurartesim treatment was compared with Artesunate + Mefloquine (AS + MQ). The primary end-point was the PCR-corrected cure rate at Day 63.

Study DM040011 was conducted in African paediatric patients with uncomplicated *P. falciparum* malaria. Eurartesim treatment was compared with Artemether + Lumefantrine (A + L). The primary end-point was PCR-corrected cure rate at Day 28.

The results for the primary endpoint in the modified intent to treat (m-ITT) populations (defined as all randomised patients who received at least one dose of the study treatment, with the exclusion of those patients lost to follow up for unknown reasons) were as follows:

Study	PCR-corrected cure rate (m-ITT)			95 % two-sided CI on the treatment difference (Eurartesim - Comparator); p-value
	Eurartesim	AS + MQ	A + L	
DM040010 (n=1087)	97.0%	95.3%	-	(-0.84, 4.19)%; p=0.161
DM040011 (n=1524)	92.7%	-	94.8%	(-4.59, 0.45)%; p=0.128

In each case the results confirmed that Eurartesim was not inferior to the comparator medicinal product. In both studies, the true treatment failure rate was below the 5% efficacy threshold set by WHO.

The age-specific PCR-corrected cure rates in the m-ITT populations are tabulated below for the Asian and African studies, respectively:

Study	PCR-corrected cure rate (m-ITT)			95% two-sided CI on the treatment difference (Eurartesim - Comparator); p-value
	Eurartesim	AS + MQ	A + L	
DM04010 (n=1087)				
≤5years	100.0%	100.0%	-	-
>5 to ≤12years	98.2%	96.5%	-	(-3.67, 7.09)%; 0.605
>12 to ≤18 years	97.3%	100.0%	-	(-6.40, 0.99)%;
>18 to ≤64 years	96.6%	94.4%	-	1.000

				(-0.98, 5.30)%; 0.146
DM04011 (n=1524)				(-12.66, -1.32)% ⁽¹⁾ ; 0.064
≤1 year	91.5%	-	98.5%	(-6.76, 2.63)%;
>1 to ≤2 years	92.6%	-	94.6%	0.413
>2 to ≤5 years	93.0%	-	94.0%	(-4.41, 2.47)%; 0.590

(1) This CI is asymptotic because the exact CI could not be computed

5.2 Pharmacokinetic properties

Pharmacokinetic profiles of arteminol and piperazine have been investigated in animal models and in different human populations (healthy volunteers, adult patients and paediatric patients).

Absorption

Arteminol is very rapidly absorbed, T_{max} being approximately 1-2 hrs after single and multiple dosing. In patients, mean C_{max} (CV%) and AUC_{INF} of arteminol (observed after the first dose of Eurartesim) were 752 (47%) ng/ml and 2,002 (45 %) ng/ml*h, respectively.

Arteminol bioavailability appears to be higher in malaria patients than in healthy volunteers, possibly because malaria *per se* has an effect on arteminol disposition. This may reflect malaria-associated impairment of hepatic function, causing an increase in arteminol bioavailability (reduction of first hepatic effect) without affecting its apparent elimination half-life, which is absorption rate limited. In healthy male volunteers under fasting conditions, mean C_{max} and AUC_{INF} of arteminol ranged between 180-252 ng/ml and 516-684 ng/ml*h, respectively.

The systemic exposure to arteminol was slightly lower following the last dose of Eurartesim (lower than after the first dose by up to 15%). arteminol pharmacokinetic parameters were found to be similar in healthy volunteers of Asian and Caucasian origin. arteminol systemic exposure on the last day of treatment was higher in females than in males, the difference being within 30%.

In healthy volunteers, arteminol exposure was increased by 43% when administered with a high fat/high calorie meal.

Piperazine, a highly lipophilic compound, is slowly absorbed. In humans, piperazine has a T_{max} of approximately 5 hours following a single and repeated dose. In patients mean (CV%) C_{max} and AUC_{0-24} (observed after the first dose of Eurartesim) were 179 (62%) ng/ml and 1,679 (47%) ng/ml*h, respectively. Due to its slow elimination, piperazine accumulates in plasma after multiple doses with an accumulation factor of approximately 3. Piperazine pharmacokinetic parameters were found to be similar in healthy volunteers of Asian and Caucasian origin. On the other hand, on the last day of Eurtartesim treatment, the piperazine maximum plasma concentration was higher in female than in male healthy volunteers, the difference being in the order of 30 to 50%.

In healthy volunteers, piperazine exposure is increased approximately 3-fold when administered with a high fat/high calorie meal. This pharmacokinetic effect is accompanied by an increased effect on prolongation of the QT interval. Accordingly, Eurartesim should be administered with water no less than 3 hours after the last food intake, and no food should be taken within 3 hours after each dose (see section 4.2).

Distribution

Both piperazine and arteminol are highly bound to human plasma proteins: the protein binding observed in *in vitro* studies was 44-93% for arteminol and >99% for piperazine. Moreover, from *in vitro* and *in vivo* data in animals, piperazine and arteminol tend to accumulate in RBC.

Arteminol was observed to have a small volume of distribution in humans (0.8 l/kg; CV 35.5%). Pharmacokinetic parameters observed for piperazine in humans indicate that this active substance has a large volume of distribution (730 l/kg; CV 37.5%).

Biotransformation

Artemimol is principally converted to α - artemimol- β -glucuronide (α - artemimol-G). Studies in human liver microsomes showed that artemimol was metabolised by the UDP-glucuronosyltransferase (UGT1A9 and UGT2B7) to α - artemimol-G with no cytochrome P450-mediated metabolism.

In vitro drug-drug interaction studies revealed that artemimol is an inhibitor of CYP1A2; therefore, there is the potential for artemimol to increase plasma concentrations of CYP1A2 substrates (see section 4.5).

In vitro metabolism studies demonstrated that piperazine is metabolised by human hepatocytes (approximately 85% of piperazine remained after 2 hours incubation at 37°C). Piperazine was mainly metabolised by CYP3A4 and to a lesser extent by CYP2C9 and CYP2C19. Piperazine was found to be an inhibitor of CYP3A4 (also in a time-dependent way) and to a lesser extent of CYP2C19, while it stimulated the activity of CYP2E1.

No effect on the metabolite profile of piperazine in human hepatocytes was observed when piperazine was co-incubated with artemimol. The piperazine major metabolites were a carboxyl acid cleavage product, and a mono-N-oxidated product.

In human studies, piperazine was found to be a mild inhibitor of CYP3A4 enzyme while potent inhibitors of CYP3A4 activity caused mild inhibition of piperazine metabolism (see section 4.5).

Elimination

The elimination half-life of artemimol is approximately 1 hour. The mean oral clearance for adult patients with malaria was 1.34 l/h/kg. The mean oral clearance was slightly higher for paediatric patients, however the differences were minor in magnitude (<20%). Artemimol is eliminated by metabolism (mainly glucuroconjugation). Its clearance was found to be slightly lower in female than in male healthy volunteers. Data regarding artemimol excretion in humans are scarce. However, it is reported in the literature that the excretion of unchanged active substance in human urine and faeces is negligible for artemisinin derivatives.

The elimination half-life of piperazine is around 22 days for adult patients and around 20 days for paediatric patients. The mean oral clearance for adult patients with malaria was 2.09 l/h/kg, while in paediatric patients was 2.43 l/h/kg. Due to its long elimination half-life, piperazine accumulates after multiple dosing.

Animal studies showed that radiolabelled piperazine is excreted by the biliary route, while urinary excretion is negligible.

Pharmacokinetics in special patient populations

No specific pharmacokinetic studies have been performed in patients with hepatic or renal insufficiency, or in elderly people.

In a paediatric pharmacokinetic study, and based on very limited sampling, minor differences were observed for artemimol pharmacokinetics between the paediatric and adult populations. The mean clearance (1.45 l/h/kg) was slightly faster in the paediatric patients than in the adult patients (1.34 l/h/kg), while the mean volume of distribution in the paediatric patients (0.705 l/kg) was lower than in the adults (0.801 l/kg).

The same comparison showed that piperazine absorption rate constant and terminal half-life in children were predominantly similar to those seen in adults. However, the apparent clearance was faster (1.30 versus 1.14 l/h/kg) and the apparent total volume of distribution was lower in the paediatric population (623 versus 730 l/kg).

5.3 Preclinical safety data

General toxicity

Literature data concerning chronic toxicity of piperazine in dogs and monkeys indicate some hepatotoxicity and mild reversible depression of total white cell and neutrophil counts.

The most important nonclinical safety findings after repeated dosing were the infiltration of macrophages with intracytoplasmic basophilic granular material consistent with phospholipidosis and degenerative lesions in numerous organs and tissues. These adverse reactions were seen in animals at exposure levels similar to clinical exposure levels, and with possible relevance to clinical use. It is not known whether these toxic effects are reversible.

Artemimol and piperazine were not genotoxic/clastogenic based on *in vitro* and *in vivo* testing.

No carcinogenicity studies have been performed.

Artemimol causes embryoletality and teratogenicity in rats and rabbits.

Piperazine did not induce malformation in rats and rabbits. In a perinatal and postnatal development study (segment III) in female rats treated with 80 mg/kg, some animals had a delay of delivery inducing mortality of the neonates. In females delivering normally the development, behaviour and growth of the surviving progeny was normal following exposure *in utero* or via milk. No reproduction toxicity studies have been performed with the combination of artemimol and piperazine.

Central nervous system (CNS) toxicity

There is potential for neurotoxicity of artemisinin derivatives in man and animals, which is strongly related to the dose, route and formulations of the different artemimol pro-drugs. In humans, the potential neurotoxicity of orally administered artemimol can be considered highly unlikely, given the rapid clearance of artemimol, and its short exposure (3 days of treatment for malaria patients). There was no evidence of artemimol-induced lesions in the specific nuclei in rats or dogs, even at lethal dose.

Cardiovascular toxicity

Effects on blood pressure and on PR and QRS duration were observed at high piperazine doses. The most important potential cardiac effect was related to cardiac conduction.

In the hERG test, the IC₅₀ was 0.15 µmol for piperazine and 7.7 µmol for artemimol. The association of artemimol and piperazine does not produce hERG inhibition greater than that of the single compounds.

Phototoxicity

There are no phototoxicity concerns with artemimol, as it does not absorb in the range of 290-700 nm.

Piperazine has an absorption maximum at 352 nm. Since piperazine is present in the skin (about 9% in the non-pigmented rat and only 3% in the pigmented rat), slight phototoxic reactions (swelling and erythema) were observed 24 hours after oral treatment in mice exposed to UV radiation.

6. Pharmaceutical particulars

6.1 List of excipients

Mannitol
Lactose
Sodium Lauryl Sulphate
Carmellose Sodium
PVP K-30
Isopropyl Alcohol
Purified Talc
Carmellose Sodium
Colloidal Anhydrous Silica (Aerosil)

Magnesium Stearate
Coating
Insta Coat Universal (IC-U-1308 -White)
Indigo Carmine Supra
Methylene dichloride
Isopropyl Alcohol

6.2 Incompatibilities

Not applicable.

6.3 Shelf life

3 years.

6.4 Special precautions for storage

Do not store above 30°C.

Store in the original package in order to protect from light and moisture.

6.5 Nature and contents of container

9 tablet packed in one Alu/Pvc blister.

6.6 Special precautions for disposal and other handling

No special requirements.

7. Applicant of Manufacturer

NAFDAC Reg. No. B4-2194

MAXHEAL LABORATORIES PVT. LTD.

PLOT NO.2-7/80-85 SURSEZ, SACHIN, DIST.

SURAT (GUJARAT) 394 230